

Two-Loop Diagrams in Noncommutative φ_4^4 theory

I. Ya. Aref'eva[§], D.M. Belov[†] and A.S. Koshelev[†]

[§] *Steklov Mathematical Institute,
Gubkin st.8, Moscow, Russia, 117966*
arefeva@genesis.mi.ras.ru

[†] *Physical Department, Moscow State University,
Moscow, Russia, 119899*
belov@orc.ru, kas@depni.npi.msu.su

Abstract

Explicit two-loop calculations in noncommutative φ_4^4 theory are presented. It is shown that the model is two-loop renormalizable.

1 Introduction

Noncommutative geometry[1] and quantum groups [2] are of relevance to quantization of space-time (see, for example, [3, 4, 5] and references therein). Another possibility of an application of noncommutative geometry deals with fields that take values on q-deformed "manifolds", in particular on quantum planes [4] or on quantum groups [6]. These two approaches are closely related. For example, gauge theory on noncommutative torus is equivalent to noncommutative gauge theory on the commutative torus. One of motivations to consider noncommutative field theories is a hope that it would be possible to cure quantum field theory divergences[7, 3, 8].

The renovation of the interest in noncommutative field theories has been stimulated by the paper of Connes, Douglas and Schwarz [9], where it was shown that supersymmetric gauge theory on noncommutative torus is naturally appeared in compactification of Matrix theory [10] (see [11] for further developments). The appearance of noncommutative geometry in string theory with a nonzero B-field [9, 12, 13] and explicit construction of a change of variables that shows an equivalence between ordinary gauge fields [13] and noncommutative gauge fields raises a question about selfconsistency of noncommutative Yang-Mills (NCYM) theory. NCYM theory is related with deformation quantization, see [14]. In such type of models ultraviolet divergences are still present [15]. Moreover, renormalizability is not

evident for noncommutative field theories. Nonrenormalizability of NCYM theory would mean inconsistency of a string theory in the B-field background at least at the zero-mode level. Therefore, it is crucial from string point of view to clarify this question. Explicit calculations performed at one-loop level show renormalizability in this approximation [16]. The next orders have not been calculated yet. About a general discussion of renormalizability see [18].

The goal of this paper is to show renormalizability of the noncommutative scalar theory in two-loop approximation. We will construct explicitly one and two-loop counterterms and show that renormalized 1PI functions are well-defined for non-exceptional momenta (compare with IR finiteness of renormalized 1PI functions in massless theories). The similar calculations for NCYM are in progress. Note that there are more similarity between calculations of counterterms in NCYM theory and those in charged scalar field theory [14].

The paper is organized as follows. In Sect.2 we formulate the model and explain why we will work in the language of single-line graphs and symmetric vertices. In Sect.3 we present the one loop calculations and in Sect.4 we schematically present two-loop calculations. Details of these calculations are collected in Appendix. In the last Section we note a problem with IR divergences for higher loops renormalized 1PI functions (we have this problem in spite of we deal with a massive theory).

2 The Model

We consider the theory with the action

$$S = S_0 + S_{int} = \int d^d x \left[\frac{1}{2} (\partial_\mu \varphi)^2 + \frac{1}{2} m^2 \varphi^2 + \frac{g}{4!} (\varphi \star \varphi \star \varphi \star \varphi)(x) \right], \quad (1)$$

where $\star$ is a Moyal product $(f \star g)(x) = e^{i\xi \theta^{\mu\nu} \partial_\mu \otimes \partial_\nu} f(x) \otimes g(x)$, ξ is a deformation parameter, $\theta^{\mu\nu}$ is a non-degenerate skew-symmetric real constant matrix, $\theta^2 = -1$, d is even. $p_1 \wedge p_2 \equiv \xi p_1 \theta p_2$. Let us rewrite the interaction in the Fourier components,

$$S_{int} = \frac{g}{4!(2\pi)^d} \int dp_1 dp_2 dp_3 dp_4 e^{-ip_1 \wedge p_2 - ip_3 \wedge p_4} \varphi(p_1) \varphi(p_2) \varphi(p_3) \varphi(p_4) \delta(p_1 + p_2 + p_3 + p_4). \quad (2)$$

We see the following distinguish features of the deformed theory as compared with standard local φ_d^4 model:

- There are non-local phase factors in the vertex.
- These factors provide regularization for some loop integrals but not for all [16, 14, 17].
- To have renormalizability the sum of divergences in each order of perturbation theory must have a phase factor already present in the action.

To single out phase factors it is convenient to use the 't Hooft double-line graphs and a notion of planar graphs. For planar graphs the phase factors do not affect the Feynman integrations at all [15]. In particular the planar graphs have exactly the same divergences as in the commutative theory[17]. There are no superficial divergences in nonplanar graphs since they are regulated by the phase factors [16, 14, 17, 18]. Moreover, oscillating phases regulate also divergent subgraphs, unless they are not divergent planar subgraphs.

So, at first sight it seems that the proof of renormalizability is rather trivial. One has divergences only in planar graphs, so one can use the fact that planar theory is renormalizable




Figure 1: Planar graph with a nonplanar subgraph

if its scalar counterpart does [20]. Or in other words, one can take the planar approximation of ordinary theory, find divergences within this approximation and write counterterms in noncommutative theory as divergent parts of planar graphs multiplied on the phase factors. However these arguments work only for superficial divergences. The situation is more subtle for divergent subgraphs. The reason is that a planar graph can contain nonplanar subgraphs (see simple example on Fig.1) and these divergences should be also removed. Therefore, renormalizability of the theory (1) is not obvious.

In explicit calculations we will use single-line graphs and symmetric vertices. After symmetrization of (2) we get

$$S_{int} = \frac{g}{3 \cdot 4!} \frac{1}{(2\pi)^d} \int dp_1 dp_2 dp_3 dp_4 \varphi(p_1) \varphi(p_2) \varphi(p_3) \varphi(p_4) \delta(p_1 + p_2 + p_3 + p_4) \times [\cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) + \cos(p_1 \wedge p_3) \cos(p_2 \wedge p_4) + \cos(p_1 \wedge p_4) \cos(p_2 \wedge p_3)] \quad (3)$$

and the vertex is a sum of three terms (see Fig.2).




Figure 2: The symmetric vertex

All further examination is confined to the case $d = 4$.

3 One Loop Counterterms

In this section we will compute explicitly one-loop counterterms using dimensional regularization $d = 4 - 2\epsilon$. We will also present the explicit form of finite part for two point and four point 1PI functions, $\Gamma^{(2)}$ and $\Gamma^{(4)}$, in the one loop approximation. We use the standard notations for perturbation expansion of 1PI-functions, $\Gamma^{(i)} = \sum_n g^n \Gamma_n^{(i)}$, $\Gamma^{(2)} = \Gamma_{f.p.}^{(2)} + \Delta\Gamma^{(2)}$ and $\Gamma^{(4)} = \Gamma_{f.p.}^{(4)} + \Delta\Gamma^{(4)}$



Figure 3: $\Gamma_1^{(2)} - \Delta\Gamma_1^{(2)}$

The only graph 3a contributes to $\Gamma_1^{(2)}$ and

$$\Gamma_1^{(2)} = -\frac{g}{6(2\pi)^d} \int dk \frac{2 + \cos 2p \wedge k}{k^2 + m^2} =$$

$$\frac{g}{32\pi^2} \frac{2}{3} \left[m^2 \left(\frac{1}{\epsilon} + \psi(2) - \ln \frac{m^2}{4\pi\mu^2} \right) - \sqrt{\frac{m^2}{\xi^2 p^2}} K_1(2m\xi|p|) \right]. \quad (4)$$

Its divergent part is subtracted by the counterterm 3b. Note that the representation (4) takes place only for $p \neq 0$.

$\Gamma_2^{(4)}$ is a sum of s-, t- and u-channel graphs, $\Gamma_2^{(4)} = \Gamma_{2,s}^{(4)} + \Gamma_{2,t}^{(4)} + \Gamma_{2,u}^{(4)}$. The explicit form of $\Gamma_{2,s}^{(4)}$ is

$$\Gamma_{2,s}^{(4)} = \frac{g^2}{18(2\pi)^d} \int \frac{\mathcal{P}_{4a}(\{p\}, k)}{(k^2 + m^2)((P + k)^2 + m^2)} dk \quad (5)$$

where $P = p_1 + p_2$, $\mathcal{P}_{4a}(\{p\}, k) = [2 \cos(p_1 \wedge p_2) \cos(k \wedge P) + \cos(p_1 \wedge p_2 + (p_1 - p_2) \wedge k)] \times [2 \cos(p_3 \wedge p_4) \cos(k \wedge P) + \cos(p_3 \wedge p_4 + (p_4 - p_3) \wedge k)]$. The trigonometric polynomial $\mathcal{P}_{4a}$ can be rewritten in the form

$$\mathcal{P}_{4a} = 2 \cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) + \sum_j ' c_j e^{i\Phi^j(p) + i b^j(p) \wedge k} \quad (6)$$

where the sum $\sum'$ goes over all j for which linear functions $b^j(p)$ are nonzero for almost all $\{p\}$. The only first term in (6) contributes to a pole part (see Fig. 4a) and we have




Figure 4: a) $\Delta\Gamma_{2,s}^{(4)}$ b) Cross denotes the 4-vertex counterterm

$$\begin{aligned} \Gamma_{2,s}^{(4)} = & \frac{g^2}{32\pi^2} \frac{2}{9} \left[\cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) \left(\frac{1}{\epsilon} + \psi(1) - \int_0^1 dx \ln \frac{m^2 + x(1-x)P^2}{4\pi\mu^2} \right) + \right. \\ & \left. + \sum_j ' c_j e^{i\Phi^j(p)} \int_0^1 dx K_0(\sqrt{\xi^2 b^j{}^2 (m^2 + P^2 x(1-x))}) e^{ix b^j \wedge P} \right]. \end{aligned} \quad (7)$$

The representation (7) is well defined only if all b^j are nonzero, i.e. for non exceptional momenta. It is the matter of simple algebra to sum up the divergent parts of $\Gamma_{2,s}^{(4)}$, $\Gamma_{2,t}^{(4)}$ and $\Gamma_{2,u}^{(4)}$ to obtain graph with symmetric vertex.

Therefore, in the one-loop we have

$$\Delta\Gamma_{1l}^{(2)} = \frac{g}{48\pi^2} \frac{m^2}{\epsilon} \quad (8)$$

$$\Delta\Gamma_{1l}^{(4)} = \frac{g^2}{16\pi^2} \frac{1}{9\epsilon} [\cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) + \cos(p_1 \wedge p_3) \cos(p_2 \wedge p_4) + \cos(p_1 \wedge p_4) \cos(p_2 \wedge p_3)] \quad (9)$$

4 Two Loop Counterterms

Let us consider $\Gamma_2^{(2)}$. The corresponding graphs are shown in Fig. 5. Graph 5a is compensated by graph 5c (see Appendix B).

Let us consider graph 5b (an explicit expression for this graph is presented in Appendix C, eq.(C.4)). This graph has superficial divergence which is removed by a local counterterm.




Figure 5: Graphs contributing to $\Gamma_2^{(2)}$

Also it has divergent subgraphs. Considering one of them (Fig 6a) ($b_{12} = 0, b_1 = 0, b_2 \neq 0$, i.e. case i) in Appendix C) we see that a contribution of this divergent subgraph can be represented as 6d which is nothing but a non-local tadpole with a one-loop renormalized coupling constant. This shows the two-loop renormalizability of $\Gamma^{(2)}$.




Figure 6: Divergent subgraph of 5b

Let us now prove two-loop renormalizability of $\Gamma^{(4)}$. We have three two-loop graphs (7a,b,c) and corresponding crossing graphs. In the same order of g there are also graphs with counterterms, (7d,e,f).




Figure 7: a), b) and c) show graphs contributing to $\Gamma_3^{(4)}$; d), e) and f) show graphs with counterterms (cross denotes counterterm (8))

Graph 7a has only one divergent subgraph and this divergence is compensated by graph 7d. More interesting is a compensation of divergences of graphs 7b and 7c caused by one-loop divergent subgraphs. As we will see these divergences are compensated by 7e only if we take the sum of graphs 7b and 7c. Indeed, we have equalities presented in fig.8. The second equality in fig.8 is due to symmetry under exchange $1' \leftrightarrow 4'$. So, we have a compensation of the terms presented in fig.7e and 8d. This compensation is checked explicitly in Appendix.

Therefore, in two loops we have

$$\Delta\Gamma_{2l}^{(2)} = \pi^4 \frac{g^2}{3^2(2\pi)^8} \left[\frac{5m^2}{8\epsilon^2} - \frac{3m^2}{4\epsilon} \left(\frac{3}{2} + \Psi(1) - \ln \frac{m^2}{4\pi\mu^2} \right) - \frac{p^2}{16\epsilon} \right] \quad (10)$$

$$\begin{aligned} \Delta\Gamma_{2l}^{(4)} = \frac{g^3}{3^3(2\pi)^8} \pi^4 \left[\frac{1}{\epsilon^2} - \frac{1}{2\epsilon} \right] & [\cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) + \\ & + \cos(p_1 \wedge p_3) \cos(p_2 \wedge p_4) + \cos(p_1 \wedge p_4) \cos(p_2 \wedge p_3)] \end{aligned} \quad (11)$$

Figure 8: Divergent parts of graphs 7b and 7c

5 Discussion and Conclusion

Let us compare our counterterms with corresponding counterterms in the ordinary theory. In one loop for $\Delta\Gamma_{1l}^{(4)}$ we have extra factor $2/9$ as compared with the local φ_4^4 theory. "2" comes from free terms of the trigonometric polynomial (6) and $1/9$ is from the vertex. We have the factor $2/9$ also for the one-loop 4-point planar graph. Relative factors of divergent parts of two-loop graphs are collected in the Table.

Graph	Noncommutative	Commutative	Planar
	$1/3$	$1/2$	$1/3$
	$1/9$	$1/2$	$1/9$
	$1/9$	$1/4$	$1/9$
	$1/36$	$1/6$	$1/36$
	$1/(27 \cdot 2)$	$1/4$	$1/(27 \cdot 2)$
	$1/(27 \cdot 2)$	$1/2$	$1/(27 \cdot 2)$

It is obvious that our renormalized 1PI functions do not admit the limit $\xi \rightarrow 0$. A behaviour of $\Gamma_1^{(2)}$ for $p^2 \rightarrow 0$ is the same as its behaviour for $\xi \rightarrow 0$. We have

$$\Gamma_{1,f.p.}^{(2)} \underset{p^2 \rightarrow 0}{\sim} \frac{c}{(\xi|p|)^2} \quad (12)$$

Caused by this asymptotic there are problems with a IR behaviour of graphs with tadpoles. They produce divergence in the IR region if the number of insertions is more then $n > 2$ (compare with an example of ref.[21] in [18]).

Perhaps, it would be also interesting to develop the theory of noncommutative quantum (gauge) fields not only on the torus but also on more general manifolds, in particular on Poisson manifolds. Perhaps the question of renormalizability of noncommutative quantum theory on certain Poisson manifolds will get a more favourable resolution [14].

Note added

After the finishing this paper we became aware of the work by Shiraz Minwalla, Mark Van Raamsdonk and Nathan Seiberg, hep-th/9912072, where mixing of the UV and the IR is discussed.

Acknowledgments

We would like to thank P. B. Medvedev, O.A. Rytchkov and I.V. Volovich for useful discussions. This work was supported in part by RFFI grant 99-01-00166 and by grant for the leading scientific schools 96-15-96208. I.A. is also supported by INTAS grant 96-0698.

Appendix

A Notations

We denote external momenta by p_i and loop momenta by k_1 and k_2 . Let $\mathcal{P}_\Gamma(\{p\}, \{k\})$ be a trigonometric polynomial corresponding to a two-loop graph Γ . It can be represented in a form

$$\mathcal{P}_\Gamma = \sum_j c_j \exp[i\Phi^j(p) + ib_1^j \wedge k_1 + ib_2^j \wedge k_2 + 2ib_{12}^j k_1 \wedge k_2], \quad (\text{A.1})$$

where $\Phi^j(p)$ are functions of external momenta, b_i^j , $i = 1, 2$ are linear functions of p_i , b_{12}^j and c_j are real numbers. We denote $\mathcal{B}_j(p_i) = c_j \exp[i\Phi^j(p)]$. Each term of (A.1) gives a contribution to α -representation of a two-loop diagram as

$$\begin{aligned} & \int e^{[-(a_1 k_1^2 + a_2 k_2^2 + 2a_{12} k_1 k_2 + l'_1 k_1 + l_2 k_2 + M^2) + ib_1 \wedge k_1 + ib_2 \wedge k_2 + 2ib_{12} k_1 \wedge k_2]} d^d k_1 d^d k_2 \prod_i d\alpha_i = \\ & \int \frac{\pi^d}{\mathcal{D}^{d/2}} \exp[-M^2 + \frac{a_1 l_2^2 + a_2 l_1^2 - 2a_{12} l_1 l_2 - \xi^2(a_1 b_2^2 + a_2 b_1^2 - 2a_{12} b_1 b_2)}{4\mathcal{D}} + \\ & i\xi \frac{a_1 l_2 \theta b_2 + a_2 l_1 \theta b_1 - a_{12}(l_1 \theta b_2 + l_2 \theta b_1)}{2\mathcal{D}} + i\xi b_{12} \frac{(l_1 - i\xi b_1 \theta) \theta (l_2 + i\xi \theta b_2)}{2\mathcal{D}}] \prod_i d\alpha_i \end{aligned} \quad (\text{A.2})$$

where $\mathcal{D} = a_1 a_2 - a_{12}^2 + \xi^2 b_{12}^2$, a_1, a_2, a_{12} depend on α_i , l_1, l_2 and M depend on α_i and p_j , b_1, b_2 depend on p_j , b_{12} is a constant.

To single out divergences we will separate the sums over j on several groups. We say that

- $j \in J(b_{12})$ if $b_{12}^j = 0$ and denote $\sum_{j \in J(b_{12})} = \sum_{b_1, b_2}$
- $j \in J(b_{12}, b_1)$ if $b_{12}^j = 0$ and $b_1 = 0$ and $b_2 \neq 0$ (for almost all p) and denote $\sum_{j \in J(b_{12}, b_1)} = \sum_{b_2}$ (similarly for $1 \leftrightarrow 2$.)
- $j \in J(b_{12}, b_1 - b_2)$ if $b_{12}^j = 0$ and $b_1 = b_2 \neq 0$ (for almost all p) and denote $\sum_{j \in J(b_{12}, b_1 - b_2)} = \sum_{b_1 = b_2}$.

B Compensation of graphs 5a and 5c

The graph 5a has the following analytic expression

$$\Gamma_{5a}(p) = \frac{1}{4} \left[\frac{g}{3(2\pi)^d} \right]^2 \int d^d k_1 d^d k_2 \frac{\mathcal{P}_{5a}(p, k_1, k_2)}{(k_1^2 + m^2)^2 (k_2^2 + m^2)},$$

$$\mathcal{P}_{5a}(p, k_1, k_2) = [2 + \cos(2k_1 \wedge k_2)][2 + \cos(2k_2 \wedge p)]. \quad (\text{B.3})$$

Considering terms with $b_{12} = 0$ in (B.3) we obtain

$$\Delta\Gamma_{5a}(p) = - \left[\frac{g}{3(2\pi)^4} \right]^2 \pi^4 \left(\frac{m^2}{\epsilon^2} + \frac{\Psi(2) + \Psi(1)}{\epsilon} + \frac{m^2}{2\epsilon} K_0(2\xi m|p|) \right) \left[\frac{4\pi\mu^2}{m^2} \right]^{2\epsilon}.$$

For graph 5c we have

$$\Gamma_{5c}(p) = \left[\frac{g}{3(2\pi)^d} \right]^2 \frac{\pi^2 m^2}{2\epsilon} \int d^d k \frac{2 + \cos(2p \wedge k)}{(k^2 + m^2)^2} = \left[\frac{g}{3(2\pi)^4} \right]^2 \pi^4 \left(\frac{m^2}{\epsilon^2} + \frac{\Psi(1)}{\epsilon} + \frac{m^2}{\epsilon} K_0(2\xi m|p|) \right) \left[\frac{4\pi\mu^2}{m^2} \right]^\epsilon.$$

We see that $\Gamma_{5a}(p) + \Gamma_{5c}(p)$ is finite.

C Two-loop corrections to $\Gamma^{(2)}(p^2)$

For Feynman graph 5b we have

$$\Gamma_{5b}(p) = \frac{1}{6} \left[\frac{g}{3(2\pi)^d} \right]^2 \int \frac{d^d k_1 d^d k_2 \mathcal{P}_{5b}(p, k_1, k_2)}{(k_1^2 + m^2)(k_2^2 + m^2)((k_1 + k_2 - p)^2 + m^2)},$$

where the trigonometric polynomial $\mathcal{P}_{5b}(p, k_1, k_2)$ has a form

$$\mathcal{P}_{5b}(p, k_1, k_2) = [1 + \cos(2k_1 \wedge k_2)][1 + \cos(2p \wedge (k_1 + k_2))] + \frac{1}{2}[1 + \cos(2k_1 \wedge k_2 + 2p \wedge (k_1 - k_2))] +$$

$$+ \cos(2p \wedge k_1) + \cos(2p \wedge k_2) + \cos(2k_1 \wedge k_2 + 2p \wedge k_1) + \cos(2k_1 \wedge k_2 - 2p \wedge k_2).$$

Using α -representation we write $\Gamma_{5b}(p)$ as

$$\Gamma_{5b}(p) = \frac{\pi^d}{6} \left[\frac{g}{3(2\pi)^d} \right]^2 \sum_j \int_0^\infty \frac{d\alpha_1 d\alpha_2 d\alpha_3 \mathcal{B}^j(p) e^{\mathcal{Q}(\alpha, b^j)}}{\mathcal{D}^{d/2}}, \quad \mathcal{D} = (\alpha_1 \alpha_2 + \alpha_2 \alpha_3 + \alpha_3 \alpha_1 + \xi^2 b_{12}^2), \quad (\text{C.4})$$

$$\mathcal{Q}(\alpha, b) = -m^2(\alpha_1 + \alpha_2 + \alpha_3) - p_1^2 \frac{\alpha_1 \alpha_2 \alpha_3 + \alpha_2 \xi^2 b_{12}^2}{\mathcal{D}} - p_1^2 \xi^2 \frac{\alpha_1 b_2^2 + \alpha_3 b_1^2 + \alpha_2 (b_1 - b_2)^2}{4\mathcal{D}}$$

Integral (C.4) may have divergences if $b_{12} = 0$ and simultaneously one of the following conditions is realized i) $b_1 = 0, b_2 \neq 0$; ii) $b_1 \neq 0, b_2 = 0$; iii) $b_1 = b_2 \neq 0$; iv) $b_1 = b_2 = 0$. We get

$$\Delta\Gamma_{5b}(p) = \frac{C_{5b}}{\epsilon^2} + \frac{C'_{5b}}{\epsilon} + \left[\frac{g}{3(2\pi)^d} \right]^2 \frac{\pi^4}{\epsilon} \sqrt{\frac{m^2}{\xi^2 p^2}} K_1(2\xi m|p|). \quad (\text{C.5})$$

For graph 5e (see also fig. 6d). we get

$$\Delta\Gamma_{5e}(p) = \frac{C_{5e}}{\epsilon^2} + \frac{C'_{5d}}{\epsilon} - \left[\frac{g}{3(2\pi)^d} \right]^2 \frac{\pi^4}{\epsilon} \sqrt{\frac{m^2}{\xi^2 p^2}} K_1(2\xi m|p|). \quad (\text{C.6})$$

Hence we see that nonlocal divergences appeared in the noncommutative theory are compensated in the sum $\Gamma_{5b}(p) + \Gamma_{5e}(p)$.

D Two-loop corrections to $\Gamma^{(4)}(p)$

The divergent part of Feynman graph 7a is exactly compensated by the graph 7d. A nontrivial compensation takes place for graphs 7b, 7c and 7e. For graph 7b we have

$$\begin{aligned} \Gamma_{7b}(p_i) = & -\frac{1}{2} \frac{g^3}{3^3(2\pi)^{2d}} \int \frac{d^d k_1 d^d k_2 \mathcal{P}_{7b}(p_i, k_1, k_2)}{(k_1^2 + m^2)(k_2^2 + m^2)((k_2 + P)^2 + m^2)((k_1 + k_2 - p_4)^2 + m^2)} = \\ & -\frac{\pi^d}{2} \frac{g^3}{3^3(2\pi)^{2d}} \sum_b \int_0^\infty \frac{d\alpha_1 d\alpha_2 d\alpha_3 d\alpha_4 \mathcal{B}^j(p_i, k_1, k_2) e^{\mathcal{Q}^j(\alpha, b^j)}}{\mathcal{D}^{d/2}} + \dots \end{aligned} \quad (\text{D.7})$$

Here dots denote the terms with $b_{12} \neq 0$ that are not relevant to our consideration,

$$\begin{aligned} \mathcal{Q}^j(\alpha, b) = & -m^2[\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4] - \frac{1}{\mathcal{D}}[P^2 \alpha_2 \alpha_3 (\alpha_1 + \alpha_4) + p_4^2 \alpha_1 \alpha_2 \alpha_4 + p_3^2 \alpha_1 \alpha_3 \alpha_4] \\ & - \frac{\xi^2}{4\mathcal{D}}[(\alpha_2 + \alpha_4)b_1^2 + \alpha_1 b_2^2 + \alpha_4(b_1 - b_2)^2] - \frac{i\xi}{\mathcal{D}}[(\alpha_1 \alpha_4 p_4 - \alpha_3(\alpha_1 + \alpha_4)P)\theta b_2 + \alpha_4(\alpha_2 p_4 - \alpha_3 p_3)\theta b_1], \end{aligned}$$

and $\mathcal{D} = (\alpha_1 \alpha_2 + \alpha_1 \alpha_3 + \alpha_1 \alpha_4 + \alpha_2 \alpha_4 + \alpha_3 \alpha_4)$. The integral in RHS of (D.7) diverges only in the cases i) $b_1 = 0, b_2 \neq 0$ and ii) $b_1 = b_2 = 0$. In the first case one gets

$$\Delta\Gamma_{7b}^{(i)} = -\frac{g^3}{3^3(2\pi)^{2d}} \frac{\pi^4}{\epsilon} \sum_{b_2} \int_0^1 dx \mathcal{B}_{b_2}(p_i) K_0 \left(\sqrt{\xi^2 b_2^2 [m^2 + P^2 x(1-x)]} \right) \exp(i\xi x b_2 \theta P). \quad (\text{D.8})$$

To determine $\mathcal{B}_{b_2}$ we pick out from $\mathcal{P}(p_i, k_1, k_2)$ a trigonometric polynomial that has $b_1 = 0$. This polynomial (after performing symmetrization $p_3 \leftrightarrow p_4$, compare with fig. 8a,b) has a form

$$\begin{aligned} \mathcal{P}_{7b}^{b_1=0}(p_i) = & [2 \cos(p_1 \wedge p_2) \cos(k_2 \wedge P) + \cos(p_1 \wedge p_2 + (p_1 - p_2) \wedge k_2)] \\ & [\cos(p_3 \wedge p_4) \cos(k_2 \wedge P) + \cos(p_3 \wedge p_4 + (p_4 - p_3) \wedge k_2)]. \end{aligned}$$

$\Delta\Gamma_{7b}^{(ii)}$ has divergences as in the local theory,

$$\Delta\Gamma_{7b}^{(ii)} = -\frac{\pi^4}{2} \frac{g^3}{3^3(2\pi)^8} \cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) \left[\frac{1}{2\epsilon^2} + \frac{1}{2\epsilon} + \frac{\Psi(1)}{\epsilon} - \frac{1}{\epsilon} \int_0^1 dx \ln \frac{m^2 + P^2 x(1-x)}{4\pi\mu^2} \right]. \quad (\text{D.9})$$

Turning to graph 7c we have

$$\Gamma_{7c}(p) = -\frac{1}{4} \frac{g^3}{3^3(2\pi)^{2d}} \int \frac{d^d k_1 d^d k_2 \mathcal{P}_{7c}(p_i, k_1, k_2)}{(k_1^2 + m^2)((P + k_1)^2 + m^2)(k_2^2 + m^2)((k_2 - P)^2 + m^2)}$$

Using the same technique as in the previous calculations we see that the graph has divergences in the cases i) $b_1 = 0, b_2 \neq 0$, ii) $b_1 \neq 0, b_2 = 0$ and iii) $b_1 = b_2 = 0$ simultaneously with $b_{12} = 0$ (in all cases). For example,

$$\Delta\Gamma_{7c}^{(ii)}(p) = -\frac{g^3}{3^3(2\pi)^2} \frac{\pi^4}{2\epsilon} \sum_{b_1} \int_0^1 dx \mathcal{B}_{b_1}(p_i) K_0 \left(\sqrt{\xi^2 b_1^2 [m^2 + P^2 x(1-x)]} \right) \exp(i\xi x b_1 \theta P). \quad (\text{D.10})$$

To determine $\mathcal{B}_{b_1}(p)$ we pick out from $\mathcal{P}_{7c}(p_i, k_1, k_2)$ a trigonometric polynomial that has $b_2 = 0$

$$\mathcal{P}_{7c}^{b_2=0} = 2 \cos(p_3 \wedge p_4) \cos(k \wedge P) [2 \cos(p_1 \wedge p_2) \cos(k \wedge P) + \cos(p_1 \wedge p_2 + (p_1 - p_2) \wedge k)].$$

$\Delta\Gamma_{7c}^{(ii)}(p)$ is obtained from (D.10) by $(1, 2) \rightarrow (3, 4)$. $\Delta\Gamma_{7c}^{(iii)}$ has divergences as in the local theory,

$$\Delta\Gamma_{7c}^{(iii)} = -\frac{\pi^4}{2} \frac{g^3}{3^3(2\pi)^8} \cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) \left[\frac{1}{\epsilon^2} + \frac{2\Psi(1)}{\epsilon} - \frac{2}{\epsilon} \int_0^1 dx \ln \frac{m^2 + P^2 x(1-x)}{4\pi\mu^2} \right]. \quad (\text{D.11})$$

Finally, graph 7e yields

$$\begin{aligned}\Gamma_{7e}(p_i) &= \frac{g^3}{3^3(2\pi)^{2d}} \frac{\pi^2}{2\epsilon} \int \frac{d^d k \mathcal{P}_{7e}(p_i, k)}{(k^2 + m^2)((k + P)^2 + m^2)} = \\ &\pi^4 \frac{g^3}{3^3(2\pi)^8} \cos(p_1 \wedge p_2) \cos(p_3 \wedge p_4) \left[\frac{1}{\epsilon^2} + \frac{\Psi(1)}{\epsilon} - \frac{1}{\epsilon} \int_0^1 dx \ln \frac{m^2 + P^2 x(1-x)}{4\pi\mu^2} \right] \\ &\frac{\pi^4}{\epsilon} \frac{g^3}{3^3(2\pi)^{2d}} \sum_b \int_0^1 dx \mathcal{B}_b(p_i) K_0 \left(\sqrt{\xi^2 b^2 [m^2 + P^2 x(1-x)]} \right) \exp(i\xi x b \theta P).\end{aligned}$$

$\sum_b$ means that the sum is over j with $b^j \neq 0$ (see (6)). The $\mathcal{B}_b(p_i)$ are coming from

$$\begin{aligned}\mathcal{P}_{7e}(p_i, k) &= [2 \cos p_1 \wedge p_2 \cos k \wedge P + \cos(p_1 \wedge p_2 + (p_1 - p_2) \wedge k)] \\ &\times [2 \cos p_3 \wedge p_4 \cos k \wedge P + \cos(p_3 \wedge p_4 + (p_4 - p_3) \wedge k)] + k].\end{aligned}$$

Due to $\mathcal{P}_{7b}^{b_1=0} + \frac{1}{2}\mathcal{P}_{7c}^{b_2=0} = \mathcal{P}_{7e}$ we get $2\Gamma_{7b} + \Gamma_{7c} + 2\Gamma_{7e} = \Delta\Gamma_{2l}^{(4)} + (finite\ terms)$.

References

- [1] A.Connes, *Noncommutative Geometry*, Academic Press, 1994.
- [2] L.D.Faddeev, N.Reshetishin and L.Taxhtajain. *Algebra and Analysis*, **1** (1987) 1.
- [3] J.Madore, "An Introduction to Noncommutative Differential Geometry and its Physical Applications", Cambridge Univ. Press, 1995.
- [4] J.Wess and B.Zumino, *Nucl.Phys.B (Proc.Suppl.)* **18B** (1990) 302.
- [5] J.Lukierski, *Deformed Quantum Relativistic Phase Spaces – an Overview*, hep-th/9812063
- [6] I.Ya.Aref'eva, and I.V.Volovich. *Quantum group gauge theory*, *Mod.Phys.Lett.* **A6** (1991) 893
- [7] I.Ya.Aref'eva and I.V.Volovich. *Quantum Group Particles and Non-Archimedean Geometry*. *Phys.Lett.* **B268** (1991) 179.
- [8] A. Kempf *Minimal Length Uncertainly Relation and Ultraviolet Regularisation*. *Phys.Rev.* **D55** (1997) 7909, hep-th/9612084
- [9] A. Connes, M.R. Douglas, A. Schwarz, "Noncommutative Geometry and Matrix Theory: Compactification on Tori", *JHEP* 02 (1998) 003.
- [10] T. Banks, W. Fischler, S.H. Shenker and L. Susskind, *Phys. Rev.***D55** (1997) 5112;
- [11] M. R. Douglas, C. Hull, "D-branes and Noncommutative Torus", *JHEP* 02 (1998) 008, M.R. Douglas, *Two Lectures on D-Geometry and Noncommutative Geometry*, hep-th/9901146.
- [12] Y.-K. E. Cheung, M. Krogh, *Noncommutative Geometry From 0-Branes in a Background B Field*, **Nucl. Phys.** **B528** (1998) 185, hep-th/9803031; Chong-Sun Chu, Pei-Ming Ho, *Noncommutative Open String and D-brane*, hep-th/9812219; M.M. Sheikh-Jabbari, *Open Strings in a B-field Background as Electric Dipoles*, hep-th/9901080; V. Schomerus, *D-Branes and Deformation Quantization*, hep-th/9903205.
- [13] N. Seiberg and E. Witten, *String Theory and Noncommutative Geometry*, *JHEP* 9909 (1999) 032, hep-th/9908142.

- [14] I.Ya.Aref'eva, and I.V.Volovich, *Noncommutative Gauge Fields on Poisson Manifolds* hep-th/9907114
- [15] T. Filk, *divergences in a field theory on quantum space*, Phys.Lett. **B376** (1996) 53-58.
- [16] C.P. Martin, D. Sanchez-Ruiz, "The one-loop UV Divergent Structure of $U(1)$ Yang-Mills Theory on Noncommutative R^4 ", Phys. Rev. Lett. **83** (1999) 476; M.M. Sheikh-Jabbari, "Renormalizability of Super Yang-Mills Theories on Noncommutative Torus", J. High Energy Phys. **9906** (1999) 015; T.Krajewski and R.Wulkenhaar, *Perturbative Quantum Gauge Fields on the Noncommutative Torus*, hep-th/9903187.
- [17] D. Bigatti and L. Susskind, *Magnetic fields, branes and noncommutative geometry*, hep-th/9908056.
- [18] I. Chepelev and R. Roiban, *Renormalization of Quantum Field Theories on Noncommutative R , I. Scalars*, hep-th/9911098
- [19] A. S. Cattaneo and G. Felder, *A path integral approach to the Kontsevich quantization formula*, math.QA/9902090
- [20] G.'t Hooft *Rigorous construction of planar diagram field theories in four dimensional Euclidean space.*, Comm.Math.Phys. **88**, 1 (1983) V. Rivasseau, *Construction and Borel Summability of Planar 4-Dimensional Euclidean Field Theory*, Comm.Math.Phys. **95**, 445-486 (1984).
- [21] N.N.Bogoliubov and D.V.Shirkov, *Introduction to the Theory of Quantum Fields*, Nauka, 1976, third edition